



# Schooner

Full Season, Medium Vine  
MIN-DAK Region



## NAVY BEAN

**Schooner** – Full Season, Medium Vine

### Profile:


Schooner is a reliable variety with a long history of consistent performance. This excellent canner has a reputation for starting strong and finishing strong, and holds its pods off the ground to avoid the costly effects of water damage. It has proven its adaptability and yielding power. Schooner germinates well in cool soils, matures evenly, and dries down uniformly.

### Economic Profile:

Assuming a 120 lbs./AC yield increase over Norstar, this variety can potentially increase your profit \$19/AC on \$16/cwt. beans minus the difference in seed cost/AC. This can calculate out to an approximate \$1,920 increased profit over Norstar on one quarter section field.



### SCHOONER VS. NORSTAR\*



\*Data from the MIN-DAK region

### TRIAL DATA

#### SCHOONER

<b>Approx. Maturity</b>	98-103 days
<b>Plant Type</b>	3B
<b>Approx. Seed Count</b>	2,804 seeds/lb.
<b>Disease Resistance</b>	<b>HR:</b> CI (Race 07) <b>IR:</b> BCMV ("1" gene)

\*See reverse side for disease resistance abbreviation chart

For customers around the world, ADM draws on its resources—its people, products, and market perspective—to help them meet today's consumer demands and envision tomorrow's needs.

[www.Seedwest.com](http://www.Seedwest.com)

800-637-5843

[seedwest@adm.com](mailto:seedwest@adm.com)





# Schooner

Full Season, Medium Vine – NAVY BEAN



## KEY TO RESISTANCE ABBREVIATIONS FOR BEANS

<b>Plant Type 1A</b>	Bush determinate erect stem
<b>Plant Type 2A</b>	Erect growth indeterminate short runners
<b>Plant Type 2B</b>	Erect growth indeterminate with medium to long runners
<b>Plant Type 3B</b>	Prostrate vine indeterminate growth with long runners
<b>BCMV</b>	Bean common mosaic caused by the specified strains of Bean common mosaic virus
<b>BCTV</b>	Curly top caused by Beet curly top virus
<b>BGYMV</b>	Bean golden yellow mosaic caused by Bean golden yellow mosaic virus
<b>CI</b>	Anthraxnose caused by <i>Collectrichum lindemuthianum</i>
<b>Psp</b>	Halo blight caused by <i>Pseudomonas savastanoi</i> pv. <i>phaseolicola</i>
<b>Pss</b>	Bacterial brown spot caused by <i>Pseudomonas syringae</i> pv. <i>syringae</i>
<b>Ua</b>	Rust caused by the specified races of <i>Uromyces appendiculatus</i>
<b>HR</b>	<b>High Resistance:</b> describes plant varieties that highly restrict the growth and development of the specified pest or pathogen under normal pest or pathogen pressure when compared to susceptible varieties. Highly resistant varieties may, however, exhibit some symptoms or damage under heavy pest or pathogen pressure.
<b>IR</b>	<b>Intermediate Resistance:</b> describes plant varieties that restrict the growth and development of the specified pest or pathogen, but may exhibit a greater range of symptoms or damage compared to highly resistant varieties. Intermediately resistant varieties will still show less severe symptoms or damage than susceptible plant varieties when grown under similar environmental conditions and/or pest or pathogen pressure.

In cases where specific races or strains are not noted the variety is resistant to some, but not necessarily all known races or strains of the pathogen.

ADM Seedwest

P.O. Box 1470

Decatur, Illinois 62525

For more information, please contact your Seedwest dry bean dealer or visit [www.Seedwest.com](http://www.Seedwest.com).

Note: All variety information presented herein is based on field and laboratory observation. Actual crop yield and quality are dependent upon many factors beyond our control and NO WARRANTY is made for crop yield and quality. Since Environmental conditions and local practices may affect variety characteristics and performance, we disclaim any legal responsibility for these. Read all tags and labels. They contain important conditions of sale, including limitations of warranties and remedies. Seedwest is a registered trademark of ADM, P.O. Box 1470, Decatur, IL 62525. [www.seedwest.com](http://www.seedwest.com)

